

Steady bubbles and drops in inviscid fluids

Lukas Niebel University of Münster

joint work with David Meyer, ICMAT Madrid
Christian Seis, University of Münster

NoKo25 — 43rd Northern German Colloquium on Applied Analysis and Numerical Mathematics

Humboldt-Universität zu Berlin, 17th October 2025

1.	Experiments	(videos,	links	on	the	last	slide)

- 2. Two-phase Euler equations with surface tension
- 3. Travelling wave solutions and an overdetermined elliptic free boundary value problem
- 4. Close-to-spherical solutions (theorem, remarks, and proof)


Two-phase Euler equations

Velocity field of the fluid $U \colon \mathbb{R} \times \mathbb{R}^3 \to \mathbb{R}^3$ solution to

$$\rho(\partial_t U + (U \cdot \nabla)U) + \nabla P = 0 \qquad \text{in } \mathbb{R} \times \mathbb{R}^3$$

$$\nabla \cdot U = 0 \qquad \text{in } \mathbb{R} \times \mathbb{R}^3$$

$$\llbracket U \cdot n \rrbracket = 0 \qquad \text{on } \mathcal{S}(t)$$



Two-phase Euler equations

Velocity field of the fluid $U: \mathbb{R} \times \mathbb{R}^3 \to \mathbb{R}^3$ solution to

$$ho(\partial_t U + (U \cdot \nabla)U) + \nabla P = 0$$
 in $\mathbb{R} \times \mathbb{R}^3$
$$\nabla \cdot U = 0$$
 in $\mathbb{R} \times \mathbb{R}^3$
$$\llbracket U \cdot n \rrbracket = 0$$
 on $\mathcal{S}(t)$

where

- $P \colon \mathbb{R} \times \mathbb{R}^3 \to \mathbb{R}$ is the pressure
- $\mathcal{S}(t)$ is the interface separating the inner $\mathcal{D}^{\sf in}(t)$ and outer $\mathcal{D}^{\sf out}(t)$ fluid domain
- $\rho(t) = \rho^{\text{in}} \mathbb{1}_{\mathcal{D}^{\text{in}}(t)} + \rho^{\text{out}} \mathbb{1}_{\mathcal{D}^{\text{out}}(t)}$ for $\rho^{\text{in}}, \rho^{\text{out}} \geq 0$, is the density function
- $[\![f]\!] = f^{\mathrm{out}} f^{\mathrm{in}}$, the jump of a quantity f across the interface.

Two-phase Euler equations

Velocity field of the fluid $U: \mathbb{R} \times \mathbb{R}^3 \to \mathbb{R}^3$ solution to

$$\rho(\partial_t U + (U \cdot \nabla)U) + \nabla P = 0 \qquad \text{in } \mathbb{R} \times \mathbb{R}^3$$

$$\nabla \cdot U = 0 \qquad \text{in } \mathbb{R} \times \mathbb{R}^3$$

$$\llbracket U \cdot n \rrbracket = 0 \qquad \text{on } \mathcal{S}(t)$$

where

- $P \colon \mathbb{R} \times \mathbb{R}^3 \to \mathbb{R}$ is the pressure
- $\mathcal{S}(t)$ is the interface separating the inner $\mathcal{D}^{\mathsf{in}}(t)$ and outer $\mathcal{D}^{\mathsf{out}}(t)$ fluid domain
- $\rho(t) = \rho^{\text{in}} \mathbb{1}_{\mathcal{D}^{\text{in}}(t)} + \rho^{\text{out}} \mathbb{1}_{\mathcal{D}^{\text{out}}(t)}$ for $\rho^{\text{in}}, \rho^{\text{out}} \geq 0$, is the density function.

Ill-posed due to Kelvin-Helmholtz instability!

Two-phase Euler equations with surface tension

Velocity field of the fluid $U: \mathbb{R} \times \mathbb{R}^3 \to \mathbb{R}^3$ solution to

$$ho(\partial_t U + (U \cdot \nabla)U) + \nabla P = 0$$
 in $\mathbb{R} \times \mathbb{R}^3$
$$\nabla \cdot U = 0$$
 in $\mathbb{R} \times \mathbb{R}^3$
$$\llbracket P \rrbracket = \sigma H$$
 on $\mathcal{S}(t)$
$$\llbracket U \cdot n \rrbracket = 0$$
 on $\mathcal{S}(t)$

where

- we take into consideration the Young-Laplace law
- H is the mean curvature (H = 2 for the unit ball)
- $\sigma>0$ is the surface tension

Two-phase Euler equations with surface tension

Velocity field of the fluid $U \colon \mathbb{R} \times \mathbb{R}^3 \to \mathbb{R}^3$ solution to

$$ho(\partial_t U + (U \cdot \nabla)U) + \nabla P = 0 \qquad \text{in } \mathbb{R} \times \mathbb{R}^3$$

$$\nabla \cdot U = 0 \qquad \text{in } \mathbb{R} \times \mathbb{R}^3$$

$$\llbracket P \rrbracket = \sigma H \qquad \text{on } \mathcal{S}(t)$$

$$\llbracket U \cdot n \rrbracket = 0 \qquad \text{on } \mathcal{S}(t)$$

Literature:

- Lots of physics literature. Influential: Hou-Lowengrub-Shelly '97
- Locally well-posed: Iguchi-Tanaka-Tani '97, Ambrose '02, Schweizer '05,

Ambrose-Masmoudi '2007, Cheng-Coutand-Shkoller '08,

- Coutand-Shkoller '08
- A priori regularity: Shatah-Zeng '08
- Finite-time singularities: Coutand-Shkoller '14,
- Castro-Córdoba-Fefferman-Gancedo-Gómez-Serrano '12

We make the ansatz

$$u(x) = U(t, x_1, x_2, x_3 + Vt) - Ve_3$$

 $p(x) = P(t, x_1, x_2, x_3 + Vt)$
 $S(t) = S + tVe_3$,

for some speed $V \ge 0$.

The time-independent u, p, \mathcal{S} solve the steady two-phase Euler equations

$$\rho(u \cdot \nabla)u + \nabla p = 0 \qquad \text{in } \mathbb{R}^3 \setminus \mathcal{S},$$

$$\nabla \cdot u = 0 \qquad \text{in } \mathbb{R}^3,$$

$$\llbracket p \rrbracket = \sigma H \qquad \text{on } \mathcal{S},$$

$$u \cdot n = 0 \qquad \text{on } \mathcal{S}.$$

with $\lim_{|x|\to\infty} u(x) = -Ve_3$.

Bernoulli equations (for steady flows) for the inner/outer phase are

$$\frac{\rho^{\mathrm{in}}}{2} \left| u^{\mathrm{in}} \right|^2 + p^{\mathrm{in}} = \mathrm{const},$$

$$\frac{\rho^{\mathrm{out}}}{2} \left| u^{\mathrm{out}} \right|^2 + p^{\mathrm{out}} = \mathrm{const}.$$

We rewrite the interfacial condition

$$[\![P]\!] = \sigma H \ \ \text{on} \ \ \mathcal{S}$$

Rewrite the interfacial condition as

$$\frac{1}{2} \llbracket \rho |u|^2 \rrbracket + \sigma H = \text{const on } \mathcal{S}.$$

We are interested in axisymmetric and swirl-free vector fields (u=u(r,z)) and azimutal component $u_{\varphi}=0$.

We assume uniform vorticity distribution in the inner phase, i.e.

$$\operatorname{curl} u^{\mathsf{in}} = \omega_{\mathsf{a}} = \frac{15}{2} a \begin{pmatrix} -x_2 \\ x_1 \\ 0 \end{pmatrix} = \frac{15}{2} a \operatorname{re}_{\varphi}$$

for $a \in \mathbb{R}$.

The fluid in the outer domain is assumed to be irrotational $\operatorname{curl} u^{\operatorname{out}} = 0$.

The volume is $|S| = \frac{4}{3}\pi R^3$.

We work with the vector stream function $\psi \colon \mathbb{R}^3 \to \mathbb{R}^3$ with

$$u = \operatorname{curl} \psi - Ve_3$$
.

The tangential flow and the axisymmetry no-swirl condition yields


$$\psi = \frac{V}{2} r e_{\varphi}$$
 on \mathcal{S} .

The identity $\operatorname{curl} \operatorname{curl} = \nabla \nabla \cdot - \Delta$ implies

$$-\Delta \psi = \omega_{\mathsf{a}} \mathbb{1}_{\mathcal{D}^{\mathsf{in}}} \text{ in } \mathbb{R}^3 \setminus \mathcal{S}.$$

The jump condition becomes

$$\frac{1}{2} \left[\left[\rho \left| \operatorname{curl} \psi - V e_3 \right|^2 \right] + \sigma H = \text{const on } \mathcal{S}.$$



Streamlines of $\psi_{\mathcal{S}}$ in axisymmetric coordinates

A first solution is given by S the sphere of radius R

$$\psi_{S}(x) = \begin{pmatrix} -x_{2} \\ x_{1} \\ 0 \end{pmatrix} \cdot \begin{cases} \frac{3a}{4} \left(R^{2} - |x|^{2} \right) + \frac{V_{S}}{2} & \text{for } |x| \leq R \\ \frac{V_{S}}{2} \frac{R^{3}}{|x|^{3}} & \text{for } |x| > R, \end{cases}$$

where $V_S = |a| R^2 \sqrt{\frac{
ho^{\rm in}}{
ho^{\rm out}}}$ is determined such that

$$\frac{1}{2} \llbracket \rho | \operatorname{curl} \psi_{S} - V_{S} e_{3} |^{2} \rrbracket = \frac{9}{8R^{2}} \left(a^{2} R^{4} \rho^{\mathsf{in}} - \rho^{\mathsf{out}} V_{S}^{2} \right) \left(x_{1}^{2} + x_{2}^{2} \right)$$

is constant on the sphere of radius R and thus


$$\frac{1}{2} \left[\left[\rho |\operatorname{curl} \psi_{S} - V_{S} e_{3}|^{2} \right] + \sigma H = 2\sigma R = \text{const.} \right]$$

A first solution is given by S the sphere of radius R

$$\psi_{S}(x) = \begin{pmatrix} -x_{2} \\ x_{1} \\ 0 \end{pmatrix} \cdot \begin{cases} \frac{3a}{4} \left(R^{2} - |x|^{2} \right) + \frac{V_{S}}{2} & \text{for } |x| \leq R \\ \frac{V_{S}}{2} \frac{R^{3}}{|x|^{3}} & \text{for } |x| > R, \end{cases}$$

with $V_S = |a| R^2 \sqrt{\frac{\rho^{\rm in}}{\rho^{\rm out}}}$.

Vortex sheet, i.e. nonzero jump of $U_S \cdot \tau$ at S, whenever $V_S \neq aR^2$.



Streamlines of $\psi_{\mathcal{S}}$ in axisymmetric coordinates

The overdetermined free boundary value problem

Given parameters $\rho^{\rm in}, \rho^{\rm out}, a, R, V$ find surface $\mathcal S$ and stream function ψ solution to

$$\begin{cases} -\Delta \psi = \frac{15}{2} a s \sin \theta e_{\varphi} \mathbb{1}_{\mathcal{D}^{\text{in}}} & \text{in } \mathbb{R}^{3} \setminus \mathcal{S} \\ \psi = \frac{V}{2} s \sin \theta e_{\varphi} & \text{on } \mathcal{S} \\ \frac{1}{2} \left[\rho \left| \text{curl } \psi - V e_{3} \right|^{2} \right] + \sigma H = \text{const} & \text{on } \mathcal{S} \end{cases}$$

vanishing at infinity.

Spherical coordinates $(s, \theta, \varphi) \in [0, \infty) \times [0, \pi) \times [0, 2\pi)$.

The overdetermined free boundary value problem

Given parameters $\rho^{\rm in}, \rho^{\rm out}, \sigma, a, R, V$ find a surface $\mathcal S$ and a stream function ψ solution to

$$\begin{cases} -\Delta \psi = \frac{15}{2} a s \sin \theta e_{\varphi} \mathbb{1}_{\mathcal{D}^{\text{in}}} & \text{in } \mathbb{R}^{3} \setminus \mathcal{S} \\ \psi = \frac{V}{2} s \sin \theta e_{\varphi} & \text{on } \mathcal{S} \\ \frac{1}{2} \left[\left[\rho \left| \text{curl } \psi - V e_{3} \right|^{2} \right] + \sigma H = \text{const} & \text{on } \mathcal{S} \end{cases} \end{cases}$$

Weber number:
$$\mathrm{We} = \frac{\rho^{\mathrm{out}} V^2 R}{\sigma}$$

Vortex Weber number:
$$\gamma = \frac{\rho^{\text{in}} a^2 R^5}{\sigma}$$

The overdetermined free boundary value problem

Rescale to R = 1 and decompose

$$\psi = \left(\mathsf{a} \psi^\mathsf{in} + \frac{\mathit{V}}{2} \mathit{s} \sin \theta \ \mathsf{e}_\varphi \right) \mathbb{1}_{\mathcal{D}^\mathsf{in}} + \mathit{V} \psi^\mathsf{out} \mathbb{1}_{\mathcal{D}^\mathsf{out}},$$

with $\psi^{\text{in}} \colon \mathcal{D}^{\text{in}} \to \mathbb{R}^3$ solution to

$$\begin{cases} -\Delta \psi^{\text{in}} = \frac{15}{2} s \sin \theta \, e_{\varphi} & \text{ in } \mathcal{D}^{\text{in}}, \\ \psi^{\text{in}} = 0 & \text{ on } \mathcal{S}, \end{cases}$$

and $\psi^{\mathrm{out}} \colon \mathcal{D}^{\mathrm{out}} \to \mathbb{R}^3$ vanishing at infinity and solving

$$egin{cases} -\Delta \psi^{\mathsf{out}} = 0 & ext{in } \mathcal{D}^{\mathsf{out}}, \ \psi^{\mathsf{out}} = rac{1}{2} s \sin heta \, e_{arphi} & ext{on } \mathcal{S}. \end{cases}$$


Jump condition: $\frac{\gamma}{2}|\text{curl }\psi^{\text{in}}|^2 - \frac{\text{We}}{2}|\text{curl }\psi^{\text{out}} - e_3|^2 + H = \text{const on }\mathcal{S}.$

Perturbation of the spherical solution

For a shape function $\eta \in H^{\beta}(\mathbb{S}^2)$ we consider

$$S_{\eta} = \left\{ (1 + \eta(x))x : x \in \mathbb{S}^2 \right\}.$$

In axisymmetric coordinates:



Perturbation of the spherical solution

For a shape function $\eta \in H^{\beta}(\mathbb{S}^2)$ we consider

$$S_{\eta} = \{(1 + \eta(x))x : x \in \mathbb{S}^2\},$$

with $\mathcal{D}_{\eta}^{\text{in}}$ and $\mathcal{D}_{\eta}^{\text{out}}$ well-defined if $\eta > -1$.

We impose

- axi-symmetry $\eta = \eta(\theta)$, and
- reflection invariance across the reference plane, $\eta(\frac{\pi}{2} \theta) = \eta(\frac{\pi}{2} + \theta)$ and write $H_{\mathrm{sym}}^{\beta}(\mathbb{S}^2)$ for that subspace.

Set
$$\mathcal{M}^{\beta}=\{\eta\in \mathsf{H}^{\beta}_{\mathrm{sym}}(\mathbb{S}^2): \left|\mathcal{D}^{\mathsf{in}}_{\eta}\right|=\frac{4}{3}\pi \text{ and } \|\eta\|_{\mathsf{H}^{\beta}}\leq c_0\} \text{ for } c_0>0 \text{ small.}$$

Perturbative ansatz

We introduce the functional $\mathcal{F} \colon \mathbb{R} \times \mathbb{R} \times \mathcal{M}^{\alpha+2} \to \mathsf{H}^{\alpha}_{\mathrm{sym}}(\mathbb{S}^2)/_{\mathrm{const}}$ as

$$\mathcal{F}(\gamma, \text{We}, \eta) = \frac{\gamma}{2} \left| (\text{curl } \psi_{\eta}^{\text{in}}) \circ \chi_{\eta} \right|^{2} - \frac{\text{We}}{2} \left| (\text{curl } \psi_{\eta}^{\text{out}}) \circ \chi_{\eta} - e_{3} \right|^{2} + H_{\eta} \circ \chi_{\eta}$$

where $\chi_{\eta} = (1 + \eta(x))x$.

Goal: find We, γ and η such that

$$\mathcal{F}(\gamma, We, \eta) = const.$$

Spherical solution:

$$\mathcal{F}(\gamma, \gamma, 0) = 2 = \text{const.}$$

Theorem (MNS '25):

Let $\beta > 2$. There exists $c_0 = c_0(\beta) > 0$ and an increasing sequence $\Gamma = (\gamma_k)_{k \in \mathbb{N}}$ of positive numbers diverging to infinity with:

(A) For any $\gamma \in [0, \infty) \setminus \Gamma$ and any We close to but different from γ , there exists a unique nontrivial (smooth) solution $\eta = \eta(\gamma, \text{We}) \in \mathcal{M}^{\beta}$ to the jump equation (1).


If $\gamma=\varepsilon\delta^{\rm in}$ and ${\rm We}=\varepsilon\delta^{\rm out}$ for two nonnegative constants $\delta^{\rm in}\neq\delta^{\rm out}$ and a small parameter ε , we have the asymptotic expansion

$$\eta_{arepsilon} = arepsilon rac{3}{32} (\delta^{\mathsf{in}} - \delta^{\mathsf{out}}) \left(3\cos^2 \theta - 1
ight) + o(arepsilon) ext{ as } arepsilon o 0.$$


(B) For any $k \in \mathbb{N}$, there exists a unique local curve $s \mapsto \gamma(s)$ passing through γ_k and there are associated nontrivial (smooth) shape functions $\eta(s) \in \mathcal{M}^{\beta}$ such that the equation (1) is solved at $(\gamma(s), \gamma(s), \eta(s))$.

Remarks on the Theorem

k	1	2	3	4	5	6
γ_k	2.20516	3.07529	3.94492	4.81679	5.69137	6.56836



Remarks on the Theorem (A)




We $> \gamma$

 $\mathrm{We} = \gamma$

 $\mathrm{We} < \gamma$

Remarks on the Theorem (B)




 $We = \gamma$

 γ_1

 γ_2

Remarks on the Theorem



Experimental observations

Droplet Motion in Purified Systems, S. Winnikow and B. T. Chao (1966)

Corollary (MNS '25):

There exist values of γ close to the bifurcation set Γ for which non-spherical steady vortex solutions with $\mathrm{We} = \gamma$ exist. In particular, for these values, the spherical vortex is non-unique.

In the one fluid setting ($\rho^{\text{in}} = \rho^{\text{out}}$) without surface tension ($\sigma = 0$) the spherical solution with Hill's vortex core is unique up to translations (Amick-Fraenkel '86).

Remarks on the Theorem

Physics literature:

- Moore '58 derives the formal asymptotics of the shape for small Weber numbers neglecting the internal motion ($\rho^{in} = 0$).
- Harper '72 explains that the inner circulation can be approximated by Hill's vortex core.
- Pozrikidis '89 provides numerical evidence of the bifurcation branch and finds approximations for γ_1, γ_2 .

Theorem (MNS '25):

Let $\beta > 2$. There exists $c_0 = c_0(\beta) > 0$ and an increasing sequence $\Gamma = (\gamma_k)_{k \in \mathbb{N}}$ of positive numbers diverging to infinity with:

(A) For any $\gamma \in [0, \infty) \setminus \Gamma$ and any We close to but different from γ , there exists a unique nontrivial (smooth) solution $\eta = \eta(\gamma, \text{We}) \in \mathcal{M}^{\beta}$ to the jump equation (1).

If $\gamma=\varepsilon\delta^{\rm in}$ and ${\rm We}=\varepsilon\delta^{\rm out}$ for two nonnegative constants $\delta^{\rm in}\neq\delta^{\rm out}$ and a small parameter ε , we have the asymptotic expansion

$$\eta_{arepsilon} = arepsilon rac{3}{32} (\delta^{\mathsf{in}} - \delta^{\mathsf{out}}) \left(3\cos^2 \theta - 1
ight) + o(arepsilon) ext{ as } arepsilon o 0.$$

(B) For any $k \in \mathbb{N}$, there exists a unique local curve $s \mapsto \gamma(s)$ passing through γ_k and there are associated nontrivial (smooth) shape functions $\eta(s) \in \mathcal{M}^{\beta}$ such that the equation (1) is solved at $(\gamma(s), \gamma(s), \eta(s))$.

Idea of the proof

Recall

$$\mathcal{M}^{\alpha+2} = \left\{ \eta \in \mathsf{H}^{\alpha+2}_{\mathrm{sym}}(\mathbb{S}^2) : \left| \mathcal{D}^{\mathsf{in}}_{\eta} \right| = \frac{4}{3}\pi \, \, \mathsf{and} \, \, \left\| \eta \right\|_{\mathsf{H}^{\alpha+2}} \leq c_0 \right\}$$

 c_0 small; $\alpha > 0$ (\leadsto Sobolev embedding; algebra property),

$$\mathcal{F}\colon \mathbb{R} imes \mathbb{R} imes \mathcal{M}^{lpha+2} o \mathsf{H}^lpha_{\mathrm{sym}}(\mathbb{S}^2)/_{\mathrm{const}}$$
,

$$\begin{split} \mathcal{F}(\gamma,\mathrm{We},\eta) &= \frac{\gamma}{2} \left| \left(\mathrm{curl} \, \psi_{\eta}^{\mathsf{in}} \right) \circ \chi_{\eta} \right|^2 - \frac{\mathrm{We}}{2} \left| \left(\mathrm{curl} \, \psi_{\eta}^{\mathsf{out}} \right) \circ \chi_{\eta} - \mathsf{e}_3 \right|^2 + \mathsf{H}_{\eta} \circ \chi_{\eta} \\ & \mathcal{F}(\gamma,\gamma,0) = \mathrm{const} \; \mathsf{for} \; \mathsf{all} \; \gamma > 0. \end{split}$$

Idea of the proof

Observe that $\mathcal F$ is Fréchet differentiable.

We calculate

$$D_{\eta}\mathcal{F}(\gamma,\gamma,\eta)|_{\eta=0}: T_0\mathcal{M}^{\alpha+2} \to H_{\mathrm{sym}}^{\alpha}(\mathbb{S}^2)/_{\mathrm{const}},$$

which turns out to be invertible precisely if $\gamma \notin \Gamma = (\gamma_k)_{k \in \mathbb{N}}$.

- If $\gamma \notin \Gamma$ we can employ the implicit function theorem to deduce part (A).
- At $\gamma \in \Gamma$ we perform a bifurcation analysis by employing the Crandall-Rabinowitz theorem to prove part (B).

The curvature term

The curvature of a graph over the sphere can be written as

$$H_{\eta} \circ \chi_{\eta} = rac{1}{1+\eta} \left(2rac{1+\eta}{\sqrt{\mathcal{g}_{\eta}}} - rac{\Delta_{\mathbb{S}^2}\eta}{\sqrt{\mathcal{g}_{\eta}}} -
abla_{\mathbb{S}^2}rac{1}{\sqrt{\mathcal{g}_{\eta}}} \cdot
abla_{\mathbb{S}^2}\eta
ight)$$

where $g_{\eta}=(1+\eta)^2+\left|
abla_{\mathbb{S}^2}\eta
ight|^2.$

We deduce

$$D_{\eta}H_{\eta}\circ\chi_{\eta}|_{\eta=0}=-(\Delta_{\mathbb{S}^2}+2\mathrm{Id}).$$

Recall that the spherical harmonics $\{Y_l^m: l \in \mathbb{N}, -l \leq m \leq l\}$ form an eigenbasis for this operator with eigenvalues (l+2)(l-1).

A first part of the proof

At $\gamma = 0$ the curvature term dominates

$$\left.\mathrm{D}_{\eta}\mathcal{F}(0,0,\eta)\right|_{\eta=0}=-\left(\Delta_{\mathbb{S}^2}+2\mathrm{Id}\right)\colon\thinspace T_0\mathcal{M}^{\alpha+2}\to H^{\alpha}_{\mathrm{sym}}(\mathbb{S}^2)/_{\mathrm{const}}.$$

Moreover, $T_0\mathcal{M}^{\alpha+2}=\left\{\eta\in\mathsf{H}^{\alpha+2}_{\mathrm{sym}}(\mathbb{S}^2):\int_{\mathbb{S}^2}\eta\mathrm{d}\sigma=0\right\}$ and

$$\mathsf{H}_{\mathrm{sym}}^\beta(\mathbb{S}^2) := \big\{ f \in \mathsf{H}^\beta(\mathbb{S}^2) : \langle f, Y_I^m \rangle = 0 \text{ if } I \text{ is odd or } m \neq 0 \big\}.$$

Hence, the operator is invertible and we can locally solve

$$\mathcal{F}\left(\varepsilon\delta^{\mathrm{in}},\varepsilon\delta^{\mathrm{out}},\eta_{\varepsilon}\right)=\mathrm{const},\;\varepsilon\in\left[0,\varepsilon_{0}\right)$$

for functions $(\eta_{\varepsilon})_{\varepsilon \in [0,\varepsilon_0)}$. Noting that

$$\left. \mathrm{D}_{arepsilon} \mathcal{F}\left(arepsilon\delta^{\mathrm{in}}, arepsilon\delta^{\mathrm{out}}, 0
ight)
ight|_{arepsilon=0} = -rac{3}{2}\sqrt{rac{\pi}{5}}(\delta^{\mathsf{in}}-\delta^{\mathsf{out}})Y_2^0(heta)$$

gives the first-order asymptotics.

The jump term

A longer calculation reveals that

$$\langle \mathrm{D}_{\eta} \mathcal{F}(\gamma, \gamma, \eta) |_{\eta=0}, \delta \eta \rangle = rac{9}{2} \gamma \sin \theta \; e_{arphi} \cdot \left(2 \mathrm{Id} - \Lambda \right) \left(\sin \theta \; \delta \eta \; e_{arphi} \right) - \left(\Delta_{\mathbb{S}^2} + 2 \mathrm{Id} \right) \delta \eta,$$

where Λ is the Dirichlet-to-Neumann map for the Laplacian on the unit ball in \mathbb{R}^3 .

Analysis of the linearisation

We write

$$egin{aligned} [\mathcal{A}(\mu)](\delta\eta) &= rac{2}{9\gamma} \langle \left. \mathrm{D}_{\eta} \mathcal{F}(\gamma,\gamma,\eta) \right|_{\eta=0}, \delta\eta
angle \ &= \sin heta \, e_{arphi} \cdot (2\mathrm{Id} - \Lambda) (\sin heta \, \delta\eta \, e_{arphi}) - \mu (\Delta_{\mathbb{S}^2} + 2\mathrm{Id}) \delta\eta, \ \mu - rac{2}{3} \end{aligned}$$

for $\mu = \frac{2}{9\gamma}$.

Finding $\mu>0$ such that $\ker\mathcal{A}(\mu)\neq\{0\}$ is equivalent to the eigenvalue problem of the symmetric and compact operator

$$\mathcal{K} = (\Delta_{\mathbb{S}^2} + 2\mathrm{Id})^{-\frac{1}{2}}\sin\theta \ e_{\varphi} \cdot (2\mathrm{Id} - \Lambda)(\sin\theta \ \left((\Delta_{\mathbb{S}^2} + 2\mathrm{Id})^{-\frac{1}{2}}\delta\eta\right) \ e_{\varphi})$$

Analysis of the linearisation

In representation via spherical harmonics

$$\delta \eta = \sum_{k=1}^{\infty} v_k Y_{2k}^0(\theta)$$

this is an infinite matrix operator in weighted sequence spaces

$$\mathrm{h}^{lpha} := \left\{ v = (v_k)_{k \in \mathbb{N}} \, : \, \|v\|_{\mathrm{h}^{lpha}}^2 := \sum_{k=1}^{\infty} k^{2lpha} v_k^2 < \infty
ight\}.$$

Analysis of the linearisation

The operator $\mathcal K$ can be written as an infinite Jacobi matrix

$$A_k = -\frac{16k^3 + 4k^2 - 8k - 1}{64k^4 + 112k^3 + 44k^2 - 7k - 3} \sim -\frac{1}{4k}$$

$$B_k = \frac{(k+1)(2k-1)(2k+1)}{(4k+3)\sqrt{64k^6+288k^5+420k^4+180k^3-69k^2-63k-10}} \sim \frac{1}{8k}$$

Lemma:

Let $\alpha \geq 0$.

- a) For any $\mu \neq 0$, the operator $\mathcal{A}(\mu)$: $h^{\alpha+2} \to h^{\alpha}$ is a symmetric Fredholm operator of index 0.
- b) For $\mu > 0$, the nullspace $N(\mathcal{A}(\mu))$ of $\mathcal{A}(\mu)$ is at most one-dimensional and $N(\mathcal{A}(\mu)) \subset \mathsf{h}^\beta$ for all $\beta \geq 0$. Moreover, $N(\mathcal{A}(\mu)) = \{0\}$ for $\mu \leq 0$.
- c) There exists a strictly decreasing sequence $(\mu_k)_{k\in\mathbb{N}}\subset\mathbb{R}^+$ with limit 0 such that $\mathcal{A}(\mu_k)$ has a 1-dimensional nullspace and $\mathcal{A}(\mu)$ is invertible if $\mu\notin\{\mu_k:k\in\mathbb{N}\}\cup\{0\}$.
- d) We have $\mu_1 \leq \frac{\sqrt{2}}{21\sqrt{5}} + \frac{\sqrt{5}}{22\sqrt{13}} + \frac{127}{2079} \approx 0.119394$.
- e) For $0 \neq v^k \in N(\mathcal{A}(\mu_k))$, we have the transversality condition

$$D_{\mu}\mathcal{A}(\mu)\big|_{\mu=\mu_k}v^k\notin R(\mathcal{A}(\mu_k)).$$

Proof of the Theorem (A)

As before, we employ the implicit function theorem to

$$(\varepsilon, \eta) \mapsto \mathcal{F} \left(\gamma + \varepsilon \delta^{\text{in}}, \gamma + \varepsilon \delta^{\text{out}}, \eta \right)$$

whenever $\gamma \notin \Gamma$ and obtain $(\eta_{\varepsilon})_{|\varepsilon| < \varepsilon_0}$ such that

$$\mathcal{F}\left(\gamma+arepsilon\delta^{\mathrm{in}},\gamma+arepsilon\delta^{\mathrm{out}},\eta_arepsilon
ight)=\mathrm{const}$$
 for all $arepsilon\in(-arepsilon_0,arepsilon_0).$

Proof of the Theorem (B)

Theorem of Crandall and Rabinowitz '71:

Let M be a smooth Banach manifold and Y be a Banach space, $I \subset \mathbb{R}$ some open interval, and $\mathcal{G}: I \times M \to Y$ be continuous. Let $w_0 \in M$. If $(1) \mathcal{G}(\lambda, w_0) = 0$ for all $\lambda \in I$.

- (2) The Fréchet derivatives $D_{\lambda}\mathcal{G}$, $D_{w}\mathcal{G}$, $D_{\lambda w}^{2}\mathcal{G}$ exist and are continuous.
- (3) There exists $\lambda^* \in I$ and $w^* \in T_{w_0}M$ such that $Y/R(D_w\mathcal{G}(\lambda^*, w_0))$ and $N(D_w\mathcal{G}(\lambda^*, w_0)) = \operatorname{span}(w^*)$ is 1-dimensional.
- $(4) D_{\lambda w}^2 \mathcal{G}(\lambda, w)|_{(\lambda, w) = (\lambda^*, w_0)} w^* \notin R(D_w \mathcal{G}(\lambda^*, w)|_{w = w_0}).$

Then there exists a continuous local bifurcation curve $\{(\lambda(s),w(s))\}_{|s|<\varepsilon}$ with ε small such that $(\lambda(0),w(0))=(\lambda^*,w_0)$ and

$$\{(\lambda, w) \in U : w \neq w_0, \mathcal{G}(\lambda, w) = 0\} = \{(\lambda(s), w(s)) : 0 < |s| < \varepsilon\}$$

for some neighbourhood U of $(\lambda^*, w_0) \in I \times M$. Moreover,

$$w(s) = w_0 + sw^* + o(s)$$
 in M , $|s| < \varepsilon$.

Proof of the Theorem (B)

We apply the theorem of Crandall-Rabinowitz to

$$\mathcal{F} \colon (0,\infty) \times \mathcal{M}^{\alpha+2} \to H^{\alpha}_{\mathrm{sym}}(\mathbb{S}^2)/_{\mathrm{const}}, \quad (\gamma,\eta) \mapsto \mathcal{F}\left(\gamma,\gamma,\eta\right).$$

As

$$D_{\eta}\mathcal{F}(\gamma,\eta)|_{\eta=0} = \frac{9}{2}\gamma\mathcal{A}\left(\frac{2}{9\gamma}\right): T_0\mathcal{M}^{\alpha+2} \to \mathsf{H}^{\alpha}_{\mathrm{sym}}(\mathbb{S}^2)/_{\mathrm{const}}$$

the assumptions (1)–(4) in the theorem of Crandall and Rabinowitz are a consequence of the previous lemma.

References



D. Meyer, L. Niebel, and C. Seis. *Steady bubbles and drops in inviscid fluids*, Calc. Var. (64)299 (2025).



lukasniebel.github.io

Video references

- 1. https://www.youtube.com/watch?v=GmiivJkfoLg
- 2. https://www.youtube.com/watch?v=NjB7LXSQoQc
- 3. https://www.youtube.com/shorts/StysjXb9isQ